

## THE PREDICTION OF PLANT GROWTH PARAMETERS TO RNN ARCHITECTURE

ADAPA NARESH, Student, M.Tech (CSE), NIMRA COLLEGE OF ENGINEERING &  
TECHNOLOGY, A.P., India.

SYED SADAT ALI ALIAS ABDUL GANI, Professor, Dept. of Computer Science &  
Engineering, NIMRA COLLEGE OF ENGINEERING & TECHNOLOGY, A.P., India.

**Abstract** — In this paper, provide crucial role indecision making for complex agricultural problems. Data visualisation is also equally important to understand the general trends of the effect of various factors influencing the crop yield. a new deep recurrent neural network (RNN), using the Long Short-Term Memory (LSTM) neuron model, in the prediction formulations. Both the former yield, growth and stem diameter values, as well as the microclimate conditions, are used by the RNN architecture to model the targeted growth parameters. A comparative study is presented, using ML methods, such as support vector regression and random forest regression, utilising the mean square error criterion, in order to evaluate the performance achieved by the different methods. Very promising results, based on data that have been obtained from two greenhouses, in Belgium and the UK, in the framework of the EU Interreg SMARTGREEN project (2017-2021), are presented.

### INTRODUCTION

The present study examines the application of data visualisation techniques to find correlations between the climatic factors and rice crop yield.

The study also applies data mining techniques to extract the knowledge from the historical agriculture data set to predict rice crop yield for season of Tropical Wet and Dry climatic zone of India. The experimental results provided include sensitivity, specificity, accuracy, F1 score, Mathew's correlation coefficient, mean absolute error, root mean squared error, relative absolute error and root relative squared error. General trends in the data visualisation show that decrease in precipitation in the selected climatic zone increases the rice crop yield and increase in minimum, average or maximum temperature for the season increases the rice crop yield. For the current data set experimental results show that J48 and LAD Tree achieved the highest accuracy, sensitivity and specificity.

### PROPOSED SYSTEM

We deploy a new deep recurrent neural network (RNN), using the Long Short-Term Memory (LSTM) neuron model, in the prediction formulations. Both the former yield, growth and stem diameter values, as well as the microclimate conditions, are used by the RNN architecture to

model the targeted growth parameters. A comparative study is presented, using ML methods, such as support vector regression and random forest regression, utilising the mean square error criterion, in order to evaluate the performance achieved by the different methods. Very promising results, based on data that have been obtained from two greenhouses, in Belgium and the UK, in the framework of the EU Interreg SMARTGREEN project (2017-2021), are presented.

In order to solve the above problem, all customers must be motivated to give a rating. This paper introduces an approach for a restaurant rating system that asks every customer for a rating after their visit to increase the number of ratings as much as possible. This system can be used unmanned restaurants; the scoring system is based on facial expression detection using pretrained convolutional neural network (CNN) models. It allows the customer to rate the food by taking or capturing a picture of his face that reflects the corresponding feelings. Compared to text-based rating system, there is much less information and no individual experience reports collected. However, this simple fast and playful rating system should give a wider range of opinions about the experiences of the customers with the restaurant concept.

Effective plant growth and yield prediction is an essential task for greenhouse growers and for agriculture in general. Developing models which

can effectively model growth and yield can help growers improve the environmental control for better production, match supply and market demand and lower costs. Recent developments in Machine Learning (ML) and, in particular, Deep Learning (DL) can provide powerful new analytical tools. The proposed study utilises ML and DL techniques to predict yield and plant growth variation across two different scenarios, tomato yield forecasting and stem growth, in controlled greenhouse environments.

As there is no staff available in unmanned restaurants, it is difficult for the restaurant management to estimate how the concept and the food is experienced by the customers. Existing rating systems, such as Google and Trip Advisor, only partially solve this problem, as they only cover a part of the customer's opinions. These rating systems are only used by a subset of the customers who rate the restaurant on independent rating platforms on their own initiative. This applies mainly to customers who experience their visit as very positive or negative.

## **LITERATURE SURVEY**

(Kanai et al., 2008). The stem diameter is an important parameter describing the growth of crop plants under abiotic stress during vegetative growth stage. Therefore, it is important to generate stem diameter growth models able to predict the response

of SDV to environmental changes and plant growth under different conditions. Many studies emphasize the need to critically review and improve SDV models for assessment of environmental impact on crop growth

(Jones et al., 1999), (Heuvelink, 1996) are considered as the main applicable dynamic growth models. Those models are dependent on physiological processes, and they represent biomass partitioning, crop growth, and yield as a function of several climate and physiological parameters.

(Abreu et al., 2000) to predict tomato yield in terms of the weight of harvested fruits. The model was developed by examining the relationship between environmental parameters in a heated greenhouses in the Southern part of France. A linear relationship between flowering rate and fruit growth was the basic assumption used in this model. However, the model performance was poor when tested in unheated plastic greenhouses in Portugal.

Another tomato yield model was proposed by Adams (Adams, 2002), based on a form of graphical simulation tool. The main objective of the model was to represent weekly fluctuations of greenhouse tomato yield in terms of fruit size and harvest rate. Hourly climate data were used to estimate the rate of growth of leaf truss and the flower production. Yield seasonal fluctuations were

generally influenced by periodic variations of solar radiation and air temperature.

According to (Qaddoum et al., 2013), there is a large number of tools that can help farmers in making decisions. These can provide yield rate prediction, suggest climate control strategies, synchronise crop production with market demands.

(Vandegehuchet et al., 2014). It has been documented that SDV is sensitive to water and nutrient conditions and is closely related to the responses of crop plants to the changes of environmental conditions

### SAMPLE RESULTS





## CONCLUSION

In this paper, A comparative study is presented, using ML methods, such as support vector regression and random forest regression, utilising the mean square error criterion, in order to evaluate the performance achieved by the different methods. Very promising results, based on data that have been obtained from two greenhouses, in Belgium and the UK, in the framework of the EU Interreg SMARTGREEN project (2017-2021), are presented.

## REFERENCES

1. Abreu, P., Meneses, J. & Gary, C. 1998, "Tompousse, a model of yield prediction for tomato crops: calibration study for unheated plastic greenhouses", *XXV International Horticultural Congress, Part 9: Computers and Automation*,

*Electronic Information in Horticulture* 519, pp. 141.

2. Adams, S. 2001, "Predicting the weekly fluctuations in glasshouse tomato yields", *IV International Symposium on Models for Plant Growth and Control in Greenhouses: Modeling for the 21st Century-Agronomic and* 593, pp. 19.

3. Atanasova, N., Todorovski, L., Džeroski, S. & Kompare, B. 2008, "Application of automated model discovery from data and expert knowledge to a real-world domain: Lake Glumsø", *Ecological Modelling*, vol. 212, no. 1-2, pp. 92-98.

4. Barandiaran, I. 1998, "The random subspace method for constructing decision forests", *IEEE Transactions on Pattern Analysis and Machine Intelligence*, vol. 20, no. 8. 5. Breiman, L. 2001, "Random forests", *Machine Learning*, vol. 45, no. 1, pp. 5-32.

6. Buhmann, M.D. 2003, *Radial basis functions: theory and implementations*, Cambridge university press.

7. Chlingaryan, A., Sukkarieh, S. & Whelan, B. 2018, "Machine learning approaches for crop yield prediction and nitrogen status estimation in precision agriculture: A review", *Computers and Electronics in Agriculture*, vol. 151, pp. 61-69.

8. Cortes, C. & Vapnik, V. 1995, "Support-vector networks", *Machine Learning*, vol. 20, no. 3, pp. 273-297.

9. Daniel, J., Andrés, P., Héctor, S., Miguel, B. & Marco, T. 2008, "A survey of artificial neural network-based modeling in agroecology" in *Soft Computing applications in industry* Springer, , pp. 247-269.

10. Duchesne, L. & Houle, D. 2011, "Modelling day-to-day stem diameter variation and annual growth of balsam fir (*Abies balsamea* (L.) Mill.) from daily climate", *Forest Ecology and Management*, vol. 262, no. 5, pp. 863-872.

11. Fan, X., Kang, M., Heuvelink, E., De Reffye, P. & Hu, B. 2015, "A knowledge-and-data-driven modeling approach for simulating plant growth: A case study on tomato growth", *Ecological Modelling*, vol. 312, pp. 363-373.

12. González-Chávez, Ma del Carmen A, Carrillo-González, R., Cuellar-Sánchez, A., Delgado-Alvarado, A., Suárez-Espinosa, J., Ríos-Leal, E., Solís-Domínguez, F.A. & Maldonado-Mendoza, I.E. 2019, "Phytoremediation assisted by mycorrhizal fungi of a Mexican defunct lead-acid battery recycling site", *Science of the Total Environment*, vol. 650, pp. 3134-3144.

13. Heuvelink, E. 1996, *Tomato growth and yield: quantitative analysis and synthesis*, Heuvelink.  
Hinckley, T.M. & Bruckerhoff, D.N. 1975, "The effects of drought on water relations and stem shrinkage of *Quercus alba*", *Canadian Journal of Botany*, vol. 53, no. 1, pp. 62-72.